

Application No. 09/724,296
Amendment dated October 9, 2003
Reply to Office Action of June 10, 2003

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1-20. Canceled

21. (Currently amended) A method for cleavage of a double-stranded DNA molecule characterized by a distorted structure, wherein said distorted structure results from ultraviolet radiation damage, a photoproduct, an abasic site, mismatched nucleotide pairing, a platinum diadduct, an intercalated molecule, an insertion deletion loop of five or fewer nucleotides or alkylation of a nucleotide or a uracil residue resulting from deamination of a cytosine residue, said method comprising the step of contacting a DNA molecule characterized by a distorted structure with a broadly specific DNA damage endonuclease selected from the group of endonucleases ~~selected from the group~~ consisting of an a ~~truncated~~ stable truncated Uvelp identified by the amino acid sequence given in SEQ ID NO:4, wherein said ~~endonuclease is purified~~ Uvelp is at least 90% pure, and a fusion protein comprising a stable truncated ~~UveP1~~ Uvelp and a heterologous sequence, wherein said fusion protein is identified by the amino acid sequence given in SEQ ID ~~NO:2~~ NO:6.
22. (Previously entered) The method of claim 21, wherein said truncated Uvelp wherein said Uvelp is at least 90% pure and has the amino acid sequence as given in SEQ ID NO:4.
23. (Previously entered) The method of claim 21, wherein said fusion protein consists of the amino acid sequence given in SEQ ID ~~NO:2~~ NO:6.

Application No. 09/724,296
Amendment dated October 9, 2003
Reply to Office Action of June 10, 2003

24. (Previously entered) The method of claim 21 wherein the insertion deletion loop is of four or fewer nucleotides.
25. (Previously entered) A method for cleavage of a double-stranded DNA molecule characterized by a distorted structure, wherein said distorted structure results from an abasic site, mismatched nucleotide pairing, a platinum diadduct, an insertion deletion loop, alkylation of a nucleotide, the presence of a uracil residue resulting from deamination of a cytosine residue, said method comprising the step of contacting a DNA molecule characterized by a distorted structure with a broadly specific DNA damage endonuclease comprising an amino acid sequence selected from the group consisting of SEQ ID NO:36; SEQ ID NO:37; SEQ ID NO:38; and SEQ ID NO:39, under conditions allowing for enzymatic activity of said endonuclease, with the proviso that when the endonuclease comprises the amino acid sequence of SEQ ID NO:38, the distorted structure does not result from mismatched nucleotides.